

REMARKS

Applicants have elected the Group I claims with traverse, as discussed in the May 26, 1999 Response. In addition, in view of the outstanding requirement Under 35 U.S.C. 121 to elect a single species for prosecution, applicants elect zinc pyrithione. The structure of pyrithione is more fully delineated at page 2, lines 21-23 of the instant specification. In addition, the pyrithione structure is shown pictorially at column 1, line 25 of U.S. Patent 2,809,971 (copy attached).

No fee is believed to be due for filing this Response. However, if there are any other fees due in connection with the filing of this response, please charge them to Wiggin & Dana's Deposit Account No. 23-1665.

Respectfully submitted

DAVID F. GAVIN, ET AL.

Dated: May 27, 1999

BY Dale Lynn Carlson
Dale Lynn Carlson
Reg. No. 28,784
Attorney for Applicants

WIGGIN & DANA
One Century Tower
New Haven, CT 06508-1832
Telephone: (203) 498-4385
Fax: (203) 782-2889

34T\$01!..DOC\12800\601\146214.01

United States Patent Office

2,809,971

Patented Oct. 15, 1957

1

2,809,971

HEAVY-METAL DERIVATIVES OF 1-HYDROXY-2-PYRIDINETHIONES AND METHOD OF PREPARING SAME


Jack Bernstein and Kathryn A. Lane, New Brunswick, N. J., assignors to Olin Mathieson Chemical Corporation, New York, N. Y., a corporation of Virginia

No Drawing. Application November 22, 1955,
Serial No. 548,545

8 Claims. (Cl. 266-270)


This application is a continuation-in-part of our parent 16
application, Serial No. 358,542, filed May 29, 1953, now
abandoned.

This invention relates to, and has for its object, the
provision of: (A) heavy-metal salts of 1-hydroxy-2(1H)-
pyridinethiones (hereinafter referred to for brevity as
1-hydroxy-2-pyridinethiones) of the general Formula I:



wherein R is hydrogen, lower alkyl, lower alkoxy or
halogen, and n is a positive integer less than five; and
(B) methods of preparing same.

The compounds of this invention can be prepared by
a method which comprises interacting a 1-hydroxy-2-
pyridinethione of the Formula I, preferably in the form
of a soluble salt thereof (e. g. an alkali metal salt or
ammonium salt thereof), with a soluble compound of the
desired heavy-metal (II) in a solvent for the reactants,
and recovering the reaction product. [Reactant I may
be in tautomeric equilibrium with the corresponding 2-
mercaptopyridine-1-oxide, i. e.,




This tautomerism will not be alluded to hereinafter, it
being understood that such tautomeric form (e. g., 2-
mercaptopyridine-1-oxide) is included when referring to
the compounds of this invention either by formula such
as I, or by a name such as 1-hydroxy-2-pyridinethione.)

Examples of utilizable compounds I include the fol-
lowing (inter alia) and soluble salts thereof: 1-hydroxy-
2-pyridinethione; 1-hydroxy-3(4, 5, or 6)-methyl-2-pyri-
dinethione; 1-hydroxy-3(4, 5, or 6)-ethyl-2-pyridine-
thione; 1-hydroxy-3(4, 5, or 6)-n-butyl-2-pyridinethione;
1-hydroxy-3(4, 5, or 6)-methoxy-2-pyridinethione; 1-hy-
droxy-3(4, 5, or 6)-ethoxy-2-pyridinethione; 1-hydroxy-
3(or 5)-bromo-2-pyridinethione; 1-hydroxy-3(or 5)-
chloro-2-pyridinethione; 1-hydroxy-4(5, or 6)-chloro-2-
pyridinethione; 1-hydroxy-4(5, or 6)-dimethyl-2-pyri-
dinethione; 1-hydroxy-2(4, 5, or 6)-ethyl-
6(3, 4 or 5)-methyl-2-pyridinethione; 1-hydroxy-4,5(or
3, 6 or 5, 6)-dimethoxy-2-pyridinethione; 1-hydroxy-3,5-
dibromo-2-pyridinethione; 1-hydroxy-3,5-dichloro-2-pyri-
dinethione; 1-hydroxy-4,5,6-trimethyl-2-pyridinethione;
and 1-hydroxy-3,4,5,6-tetramethyl-2-pyridinethione. The
unsubstituted 1-hydroxy-2-pyridinethione and the mono-
substituted 1-hydroxy-2-pyridinethione are known com-

2

pounds which can be prepared as disclosed in J. A. C. S.
72, 4362 (1950). The polysubstituted 1-hydroxy-2-pyri-
dinethiones are new compounds which can be prepared
by a method comprising reacting the desired, 2-amino-
polysubstituted-pyridine with hydrobromic acid and bro-
ming and reacting the product thus formed with nitrous
acid to yield the corresponding 2-bromo-polysubstituted-
pyridine, introducing the 1-oxide group by treatment with
a peracid, such as perbenzoic or peracetic acid to form
the corresponding 2-bromo-polysubstituted-pyridine-1-
oxide of the Formula III:



wherein R' is lower alkyl, lower alkoxy, or halogen and
m is a positive integer from two through four.

To prepare the 1-hydroxy-polysubstituted-2-pyridine-
thiones, utilizable as starting materials in the preparation
of the heavy-metal salts of this invention, the compounds
of Formula III are reacted with an alkali metal sulfide,
an alkali metal hydrosulfide or preferably thiourea. If
thiourea is used, the reaction is optimally conducted in an
organic solvent, such as alcohol, at an elevated tempera-
ture, and the resulting product is converted to its alkali
metal or ammonium salt by alkalization with a base
such as an alkali metal hydroxide or ammonium hydrox-
ide or a salt of the alkali metal with a weak acid (e. g.
sodium carbonate). The reaction yields an alkali metal
or ammonium salt of a compound of Formula I, wherein
n is 2, 3, or 4, depending on the degree of substitution
of the starting pyridine reactant.

Utilizable heavy-metal compound reactants (II) utiliz-
able in the conversion of compounds of Formula I (or
their alkali metal or ammonium salts) to the heavy-metal
salts of this invention include salts in which the heavy-
metal group is (inter alia) copper, iron, manganese, tin,
mercury, cobalt, chromium, lead, gold, cadmium, nickel,
silver, zinc, titanium, arsenic, antimony, and bismuth
(the term heavy-metal, as employed herein, including
the heavy non-metals of metallic character, such as
arsenic). The reactants II may be (inter alia) nitrates,
acetates, sulfates, and halides. When a semi-metal, such
as arsenic, antimony, or bismuth, is used, in some in-
stances less than the total valence of the element is used
in the making of the heavy-metal salt, so that a basic
salt is produced as illustrated in Example 9, hereinafter.

The following examples are illustrative of, but not
limitative of, the invention:

EXAMPLE 1

Manganese salt of 1-hydroxy-2-pyridinethione

A solution of 0.99 g. (0.005 mole) of manganese
chloride tetrahydrate in 50 cc. of water is added to a
solution of 1.27 g. (0.01 mole) of 1-hydroxy-2-pyridine-
thione in 10 cc. of N sodium hydroxide. The product,
a yellow solid, precipitates immediately, and is filtered,
washed with water, alcohol and ether, and is air-dried.
Weight about 1.3 g.

EXAMPLE 2

Nickel salt of 1-hydroxy-2-pyridinethione

A solution of 1.18 g. (0.005 mole) of nickel chloride
hexahydrate in 50 cc. of water is added to a solution of

BEST AVAILABLE COPY